

Higher order generalized geometric polynomials

Levent Kargin* and Bayram Çekim**

* Alanya Alaaddin Keykubat University

Akseki Vocational School TR-07630 Antalya Turkey

Gazi University, Faculty of Science,

Department of Mathematics, Teknikokullar TR-06500,

Ankara, Turkey.

leventkargin48@gmail.com.tr and **bayramcekim@gazi.edu.tr

January 5, 2017

Abstract

According to generalized Mellin derivative [25, Eq. (2.5)], we introduce a new family of polynomials called higher order generalized geometric polynomials. We obtain some properties of them. We discuss their connections to degenerate Bernoulli and Euler polynomials. Furthermore, we find new formulas for the Carlitz's [9, Eq. (5.4)] and Howard's [23, Eq. (4.3)] finite sums. Finally, we evaluate several series in closed forms, one of which has the coefficients include values of the Riemann zeta function. Moreover, we calculate some integrals in terms of generalized geometric polynomials.

2000 Mathematics Subject Classification: 11B68, 11B83, 11M35, 33B99.

Key words: Generalized geometric polynomials, Bernoulli polynomials, Euler polynomials, Riemann zeta function.

1 Introduction

The operator $(x \frac{d}{dx})^n$, called Mellin derivative [6], has a long mathematical history. As far back as 1740, Euler used the operator as a tool work in his work [19]. The Mellin derivative and its generalizations are used to obtain a new class of polynomials [5, 17, 18, 25, 32], to evaluate some power series in closed forms [5, 7, 17, 18, 25, 27, 32] and to calculate some integrals [6, 7]. One of the generalizations of Mellin derivative is

$$\left(\beta x^{1-\alpha/\beta} D\right)^n \left[x^{r/\beta} f(x)\right] = x^{(r-n\alpha)/\beta} \sum_{k=0}^n S(n, k; \alpha, \beta, r) \beta^k x^k f^{(k)}(x), \quad (1)$$

where f is any n -times differentiable function and $S(n, k; \alpha, \beta, r)$ are generalized Stirling number pair with three free parameters (see Section 2). Stirling numbers

and their generalizations have many interesting combinatorial interpretations. Besides, these numbers are connected with some well known special polynomials and numbers [10, 11, 16, 22, 29, 28, 30, 31, 33, 37, 38]. For example, the following interesting formulas for Bernoulli numbers B_n and Euler polynomials $E_n(x)$ appear in [20]: For all $n \geq 0$;

$$B_n = \sum_{k=0}^n (-1)^k \frac{k!}{k+1} \left\{ \begin{matrix} n \\ k \end{matrix} \right\}, \quad E_n(0) = \sum_{k=0}^n (-1)^k \frac{k!}{2^k} \left\{ \begin{matrix} n \\ k \end{matrix} \right\}. \quad (2)$$

From all these motivations, by using the generalized of Mellin derivative in (1), we introduce a new family of polynomials $w_n^{(s+1)}(x; \alpha, \beta, r)$, called higher order generalized geometric polynomials, evaluate some power series in closed forms and calculate some integrals. After that, in view of the properties of higher order generalized geometric polynomials, we derive new explicit formulas for degenerate Bernoulli polynomials, Bernoulli polynomials and degenerate Euler polynomials. As a consequences of these formulas we evaluate Carlitz's [9, Eq. (5.4)] and Howard's [23, Eq. (4.3)] sums.

The summary by sections is as follows: Section 2 is the preliminary section where we give definitions and known results needed. In Section 3, we define higher order generalized geometric polynomials and obtain some properties such as recurrence relation and generating function of $w_n^{(s+1)}(x; \alpha, \beta, r)$. Moreover, we derive new explicit formulas for degenerate Bernoulli polynomials, Bernoulli polynomials and degenerate Euler polynomials. By the help of these formulas we find new formulas for Carlitz's and Howard's sums. In final section, we obtain some integral representation of $w_n^{(s+1)}(x; \alpha, \beta, r)$. Besides, we evaluate several power series in closed forms. For example, in the special cases, the following series, where the coefficients include values of the Riemann zeta function, is evaluated in closed form (for any integer $n \geq 0$)

$$\sum_{k=2}^{\infty} \frac{\zeta(k) k^n}{2^k} = \log 2 + \sum_{k=1}^n \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} k! (1 - 2^{-k-1}) \zeta(k+1).$$

Throughout this paper we assume that α, β and r are real or complex numbers.

2 Preliminaries

The generalized Stirling numbers of the first kind $S_1(n, k; \alpha, \beta, r)$ and of the second kind $S_2(n, k; \alpha, \beta, r)$ for non-negative integer m and real or complex parameters α, β and r , with $(\alpha, \beta, r) \neq (0, 0, 0)$ are defined by means of the

generating function [24]

$$\begin{aligned} \left(\frac{(1+\beta t)^{\alpha/\beta} - 1}{\alpha} \right)^k (1+\beta t)^{r/\beta} &= k! \sum_{n=0}^{\infty} S_1(n, k; \alpha, \beta, r) \frac{t^n}{n!}, \\ \left(\frac{(1+\alpha t)^{\beta/\alpha} - 1}{\beta} \right)^k (1+\alpha t)^{-r/\alpha} &= k! \sum_{n=0}^{\infty} S_2(n, k; \alpha, \beta, r) \frac{t^n}{n!}, \end{aligned}$$

with the convention $S_1(n, k; \alpha, \beta, r) = S_2(n, k; \alpha, \beta, r) = 0$ when $k > n$.

As Hsu and Shiue pointed out, the definitions or generating functions generalize various Stirling-type numbers studied previously, such as:

- i. $\{S_1(n, k; 0, 1, 0), S_2(n, k; 0, 1, 0)\} = \{s(n, k), S(n, k)\} = \left\{(-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}, \begin{Bmatrix} n \\ k \end{Bmatrix}\right\}$
are the Stirling numbers of both kinds [20, Chapter 6].
- ii. $\{S_1(n, k; \alpha, 1, -r), S_2(n, k; \alpha, 1, -r)\} = \left\{(-1)^{n-k} S_1(n, k, r + \alpha | \alpha), S_2(n, k, r | \alpha)\right\}$
are the Howard degenerate weighted Stirling numbers of both kinds [21].
- iii. $\{S_1(n, k; \alpha, 1, 0), S_2(n, k; \alpha, 1, 0)\} = \left\{(-1)^{n-k} S_1(n, k | \alpha), S_2(n, k | \alpha)\right\}$
are the Carlitz's degenerate Stirling numbers of both kinds [9].
- iv. $\{S_1(n, k; 0, -1, r), S_2(n, k; 0, 1, r)\} = \left\{(-1)^{n-k} \begin{bmatrix} n+r \\ k+r \end{bmatrix}_r, \begin{Bmatrix} n+r \\ k+r \end{Bmatrix}_r\right\}$
are the r -Stirling numbers of both kinds [8].
- v. $\{S_1(n, k; 0, \beta, -1), S_2(n, k; 0, \beta, 1)\} = \{w_\beta(n, k), W_\beta(n, k)\}$
are the Whitney numbers of the first and second kind [4].
- vi. $\{S_1(n, k; 0, \beta, -r), S_2(n, k; 0, \beta, r)\} = \{w_{\beta, r}(n, k), W_{\beta, r}(n, k)\}$
are the r -Whitney numbers of the first and second kind [31],

and so on.

According to the generalization of Stirling numbers, Hsu and Shiue [24] defined generalized exponential polynomials $S_n(x)$ as follows

$$S_n(x) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) x^k. \quad (3)$$

Later, Kargin and Corcino [25] gave an equivalent definition for $S_n(x)$ as

$$S_n(x) = \left[(x^{n\alpha-r} e^{-x}) \right]^{1/\beta} \left(\beta x^{1-\alpha/\beta} D \right)^n \left[(x^r e^x)^{1/\beta} \right], \quad (4)$$

and using the series

$$e^{x/\beta} = \sum_{k=0}^{\infty} \frac{x^k}{\beta^k k!},$$

in (4), they obtained the general Dobinski-type formula [24]

$$e^{x/\beta} S_n(x) = \sum_{k=0}^{\infty} \frac{(k\beta + r | \alpha)_n x^k}{\beta^k k!}. \quad (5)$$

Here, $(z | \alpha)_n$ is called the generalized factorial of z with increment α , defined by $(z | \alpha)_n = z(z - \alpha) \cdots (z - n\alpha + \alpha)$ for $n = 1, 2, \dots$, and $(z | \alpha)_0 = 1$. In particular, we have $(z | 1)_n = (z)_n$.

Some other properties of $S_n(x)$ can be found in [14, 15, 24, 36].

The generalized geometric polynomials $w_n(x; \alpha, \beta, r)$ are defined by means of the generalized Mellin derivative as

$$\left(\beta x^{1-\alpha/\beta} D \right)^n \left[\frac{x^{r/\beta}}{1-x} \right] = \frac{x^{(r-n\alpha)/\beta}}{1-x} w_n \left(\frac{x}{1-x}; \alpha, \beta, r \right).$$

These polynomials have the explicit formula

$$w_n(x; \alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) \beta^k k! x^k, \quad (6)$$

and the generating function

$$\sum_{n=0}^{\infty} w_n(x; \alpha, \beta, r) \frac{t^n}{n!} = \frac{(1 + \alpha t)^{r/\alpha}}{1 - x \left((1 + \alpha t)^{\beta/\alpha} - 1 \right)}, \quad \alpha\beta \neq 0. \quad (7)$$

See [25] for details.

The higher order degenerate Euler polynomials are defined by means of the generating function [9]

$$\sum_{n=0}^{\infty} \mathcal{E}_n^{(s)}(\alpha; x) \frac{t^n}{n!} = \left(\frac{2}{(1 + \alpha t)^{1/\alpha} + 1} \right)^s (1 + \alpha t)^{x/\alpha}. \quad (8)$$

From (8), we have

$$\lim_{\alpha \rightarrow 0} \mathcal{E}_n^{(s)}(\alpha; x) = E_n^{(s)}(x),$$

where $E_n^{(s)}(x)$ are the higher order Euler polynomials which are defined by the generating function

$$\sum_{n=0}^{\infty} E_n^{(s)}(x) \frac{t^n}{n!} = \left(\frac{2}{e^t + 1} \right)^s e^{xt}. \quad (9)$$

In special cases,

$$\mathcal{E}_n^{(1)}(\alpha; x) = \mathcal{E}_n(\alpha; x), \quad E_n^{(1)}(x) = E_n(x),$$

where $\mathcal{E}_n(\alpha; x)$ and $E_n(x)$ are the degenerate Euler and Euler polynomials, respectively.

The degenerate Bernoulli polynomials of the second kind are defined by means of the generating function [26]

$$\sum_{n=0}^{\infty} B_n(x | \alpha) \frac{t^n}{n!} = \frac{\frac{1}{\alpha} \log(1 + \alpha t)}{(1 + \alpha t)^{1/\alpha} - 1} (1 + \alpha t)^{x/\alpha}. \quad (10)$$

Indeed, we get

$$\lim_{\alpha \rightarrow 0} B_n(x | \alpha) = B_n(x),$$

where $B_n(x)$ are the Bernoulli polynomials which are defined by the generating function

$$\sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} = \frac{t}{e^t - 1} e^{xt},$$

with $B_n(0) = B_n$ are n -the Bernoulli numbers.

Finally, we want to mention Carlitz's degenerate Bernoulli polynomials defined by means of the generating function [9]

$$\sum_{n=0}^{\infty} \beta_n(\alpha, x) \frac{t^n}{n!} = \frac{t}{(1 + \alpha t)^{1/\alpha} - 1} (1 + \alpha t)^{x/\alpha},$$

with the relation

$$\lim_{\alpha \rightarrow 0} \beta_n(\alpha, x) = B_n(x)$$

and with $\beta_n(\alpha, 0) = \beta_n(\alpha)$ are n -the degenerate Bernoulli numbers.

3 Higher order generalized geometric polynomials

In this section, the definition of higher order generalized geometric polynomials and some properties are given. Besides, new explicit formulas for degenerate Bernoulli and Euler polynomials are derived. Some special cases of these results are studied.

For every $s \geq 0$, taking $f(x) = 1/(1-x)^{s+1}$ in (1) and using

$$\frac{\partial^k}{\partial x^k} f(x) = \frac{(s+1)(s+2)\dots(s+k)}{(1-x)^{s+k+1}},$$

we have

$$\left(\beta x^{1-\alpha/\beta} D\right)^n \left[\frac{x^{r/\beta}}{(1-x)^{s+1}} \right] = \frac{x^{(r-n\alpha)/\beta}}{(1-x)^{s+1}} \sum_{k=0}^n S(n, k; \alpha, \beta, r) \binom{s+k}{k} k! \beta^k \left(\frac{x}{1-x} \right)^k.$$

If we define the polynomials $w_n^{(s+1)}(x; \alpha, \beta, r)$ by

$$w_n^{(s+1)}(x; \alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) \binom{s+k}{k} k! \beta^k x^k, \quad (11)$$

we find

$$\left(\beta x^{1-\alpha/\beta} D \right)^n \left[\frac{x^{r/\beta}}{(1-x)^{s+1}} \right] = \frac{x^{(r-n\alpha)/\beta}}{(1-x)^{s+1}} w_n^{(s+1)} \left(\frac{x}{1-x}; \alpha, \beta, r \right). \quad (12)$$

Note that if $(s, \alpha, \beta, r) = (0, \alpha, \beta, r)$, we obtain generalized geometric polynomials [25], if $(s, \alpha, \beta, r) = (s, 0, 1, 0)$, we have general geometric polynomials [5], if $(s, \alpha, \beta, r) = (0, 0, 1, 0)$, we obtain geometric polynomials [5], if $(s, \alpha, \beta, r) = (0, 0, \beta, 1)$, we have Tanny-Dowling polynomials [4].

For $x = 1$ and $s = 0$ in (11) gives the numbers

$$w_n^{(1)}(1; \alpha, \beta, r) = B_n(\alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) \beta^k k! x^k,$$

which was defined by Corcino et al. [13]. The combinatorial interpretation and some other properties can be found in [13, 14]. Moreover, $w_n^{(s+1)}(x; \alpha, \beta, r)$ reduce to barred preferential arrangement numbers $r_{n,s}$ defined by [1, 5]

$$w_n^{(s+1)}(1; 0, 1, 0) = r_{n,s} = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \binom{s+k}{k} k!,$$

which have interesting combinatorial meaning. Therefore, we may call generalized barred preferential arrangement number pair with three free parameters for

$$B_n^{(s+1)}(\alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) \binom{s+k}{k} k! \beta^k x^k.$$

The combinatorial interpretation of these numbers may also be studied.

On the other hand, we can write $w_n^{(s+1)}(x; \alpha, \beta, r)$ in the form

$$w_n^{(s+1)}(x; \alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) \langle s+1 \rangle_k \beta^k x^k,$$

since

$$\frac{\langle x \rangle_n}{n!} = \binom{x+n-1}{n},$$

where $\langle x \rangle_n$ is the rising factorial defined by $\langle x \rangle_n = x(x+1) \cdots (x+n-1)$, for $n = 1, 2, \dots$, with $\langle x \rangle_0 = 1$. Furthermore, using the relation

$$\langle -x \rangle_n = (-1)^n \langle x \rangle_n, \quad (13)$$

if we define $w_n^{(-s)}(x; \alpha, \beta, r)$ for every $s > 0$ as

$$w_n^{(-s)}(x; \alpha, \beta, r) = \sum_{k=0}^n S(n, k; \alpha, \beta, r) (s)_k (-\beta)^k x^k,$$

we find

$$\left(\beta x^{1-\alpha/\beta} D \right)^n \left[x^{r/\beta} (1-x)^s \right] = x^{(r-n\alpha)/\beta} (1-x)^s w_n^{(-s)} \left(\frac{x}{1-x}; \alpha, \beta, r \right) \quad (14)$$

by taking $f(x) = (1-x)^s$ in (1).

We turn over $w_n^{(-s)}(x; \alpha, \beta, r)$ in Section 4.

Now, we want to deal with the properties of $w_n^{(s)}(x; \alpha, \beta, r)$. The exponential and higher order generalized geometric polynomials are connected by the relation (for any non-negative integer n and every $s > 0$)

$$w_n^{(s)}(x; \alpha, \beta, r) = \frac{1}{\Gamma(s)} \int_0^\infty z^{s-1} S_n(x\beta z) e^{-z} dz, \quad (15)$$

which is verified immediately by using (3). So, we can derive the properties of $w_n^{(s)}(x; \alpha, \beta, r)$ from those of $S_n(x)$. For example, if we use the extension of Spivey's Bell number formula to $S_n(x)$ [25, 36]

$$S_{n+m}(x) = \sum_{k=0}^n \sum_{j=0}^m \binom{n}{k} S(m, j; \alpha, \beta, r) (j\beta - m\alpha | \alpha)_{n-k} S_k(x) x^j$$

in (15) and evaluate the the integral, we derive a recurrence relation as

$$\begin{aligned} & w_{n+m}^{(s)}(x; \alpha, \beta, r) \\ &= \sum_{k=0}^n \sum_{j=0}^m \binom{n}{k} S(m, j; \alpha, \beta, r) (j\beta - m\alpha | \alpha)_{n-k} \langle s+1 \rangle_j \beta^j w_k^{(s+j)}(x; \alpha, \beta, r) x^j. \end{aligned}$$

As another application of (15), we give the following theorem.

Theorem 1 *The exponential generating function for $w_n^{(s)}(x; \alpha, \beta, r)$ is*

$$\sum_{n=0}^{\infty} w_n^{(s)}(x; \alpha, \beta, r) \frac{t^n}{n!} = \left(\frac{1}{1 - x \left((1 + \alpha t)^{\beta/\alpha} - 1 \right)} \right)^s (1 + \alpha t)^{r/\alpha}, \quad (16)$$

where $\alpha\beta \neq 0$.

Proof. From [24, Eq. (12)], let us write the generating function for $S_n(x)$ in the form

$$\sum_{n=0}^{\infty} S_n(x\beta z) \frac{t^n}{n!} = (1 + \alpha t)^{r/\alpha} \exp \left[xz \left((1 + \alpha t)^{\beta/\alpha} - 1 \right) \right].$$

Then multiply both sides by $z^{s-1}e^{-z}$ and integrate for z from zero to infinity. In view of (15) this gives

$$\sum_{n=0}^{\infty} w_n^{(s)}(x; \alpha, \beta, r) \frac{t^n}{n!} = \frac{(1 + \alpha t)^{r/\alpha}}{\Gamma(s)} \int_0^{\infty} z^{s-1} e^{-z(1-x((1+\alpha t)^{\beta/\alpha}-1))} dz.$$

Calculating the integral on the right-hand side completes the proof. ■

Setting $x = -1$ in (16), we have

$$w_n^{(s)}(-1; \alpha, \beta, r) = w_n(-1; \alpha, \beta, r) = (r - \beta s \mid \alpha)_n.$$

Some other properties of $w_n^{(s)}(x; \alpha, \beta, r)$ can be derived from (16).

Now, we attend to the connection of $w_n^{(s)}(x; \alpha, \beta, r)$ with some degenerate special polynomials.

If we take $x = -1/2$ in (16) for $\beta = 1$ and compare with (8), we have

$$w_n^{(s)}\left(\frac{-1}{2}; \alpha, 1, r\right) = \mathcal{E}_n^{(s)}(\alpha; r).$$

Thus, using (11) yields an explicit formula for higher order degenerate Euler polynomials in the following corollary.

Corollary 2 *For every $s \geq 0$, we have*

$$\mathcal{E}_n^{(s+1)}(\alpha; r) = \sum_{k=0}^n S_2(n, k, r \mid \alpha) \frac{(-1)^k \langle s+1 \rangle_k}{2^k}. \quad (17)$$

When $s = 0$, this becomes

$$\mathcal{E}_n(\alpha; r) = \sum_{k=0}^n S_2(n, k, r \mid \alpha) \frac{(-1)^k k!}{2^k}. \quad (18)$$

To the writer's knowledge, this is a new result. (17) and (18) are new results. From (16) and (9), one can obtain

$$\lim_{\alpha \rightarrow 0} w_n^{(s)}\left(\frac{-1}{2}; \alpha, \beta, r\right) = E_n^{(s)}\left(\frac{r}{\beta}\right) \beta^n.$$

Thus, we have

$$E_n^{(s)}\left(\frac{r}{\beta}\right) = \sum_{k=0}^n W_{\beta, r}(n, k) \frac{(-1)^k \langle s \rangle_k}{\beta^{n-k} 2^k},$$

which was given in [33] with a different proof. Moreover, for $\beta = 1$ and $r = 0$ the above equation reduce to the right part of (2).

Secondly, if we integrate both sides of (7) for x from -1 to 0 , we have

$$\sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n(x; \alpha, \beta, r) dx = \frac{\frac{\beta}{\alpha} \log(1 + \alpha t)}{(1 + \alpha t)^{\beta/\alpha} - 1} (1 + \alpha t)^{r/\alpha}. \quad (19)$$

In view of (10), for $\beta = 1$, the above equation becomes

$$\sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n(x; \alpha, 1, r) dx = \sum_{n=0}^{\infty} B_n(r | \alpha) \frac{t^n}{n!}.$$

Comparing the coefficients of $\frac{t^n}{n!}$ gives

$$\int_{-1}^0 w_n(x; \alpha, 1, r) dx = B_n(r | \alpha).$$

Finally, using (6) yields the following theorem.

Theorem 3 *The following equation holds for degenerate Bernoulli polynomials of the second kind*

$$B_n(r | \alpha) = \sum_{k=0}^n S_2(n, k, r | \alpha) \frac{(-1)^k k!}{k+1}.$$

We note that for $\alpha \rightarrow 0$, (19) can be written as

$$\begin{aligned} \lim_{\alpha \rightarrow 0} \sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n(x; \alpha, \beta, r) dx &= \frac{\beta t}{e^{\beta t} - 1} e^{rt} \\ &= \sum_{n=0}^{\infty} B_n\left(\frac{r}{\beta}\right) \beta^n \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{n!}$ in the above equation, we obtain

$$\lim_{\alpha \rightarrow 0} \int_{-1}^0 w_n(x; \alpha, \beta, r) dx = B_n\left(\frac{r}{\beta}\right) \beta^n.$$

Since $S_2(n, k; 0, \beta, r) = W_{\beta, r}(n, k)$ we find

$$B_n\left(\frac{r}{\beta}\right) = \sum_{k=0}^n W_{\beta, r}(n, k) \frac{(-1)^k k!}{\beta^{n-k} (k+1)},$$

which was also given in [33] with a different proof. Moreover, for $\beta = 1$ and $r = 0$ the above equation reduce to the left part of (2).

Now, we give a new explicit formula for Carlitz's degenerate Bernoulli polynomials in the following theorem.

Theorem 4 *For every $s \geq 0$,*

$$\beta_{n+1}(\alpha, r) - \beta_{n+1}(\alpha, r - s) = (n+1) \sum_{k=0}^n S_2(n, k, r | \alpha) \frac{(-1)^k \langle s \rangle_{k+1}}{k+1}. \quad (20)$$

Proof. If we integrate both sides of (16) for x from -1 to 0 , we have

$$\sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n^{(s+1)}(x; \alpha, \beta, r) dx = \frac{1}{st} \left[\frac{t(1+\alpha t)^{r/\alpha}}{(1+\alpha t)^{\beta/\alpha} - 1} - \frac{t(1+\alpha t)^{(r-\beta s)/\alpha}}{(1+\alpha t)^{\beta/\alpha} - 1} \right]. \quad (21)$$

For $\beta = 1$, the above equation becomes

$$\sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n^{(s+1)}(x; \alpha, 1, r) dx = \frac{1}{s} \left[\sum_{n=0}^{\infty} \frac{\beta_{n+1}(\alpha, r) - \beta_{n+1}(\alpha, r-s)}{n+1} \right] \frac{t^n}{n!}.$$

Equating the coefficients of $\frac{t^n}{n!}$ in the above equation, we obtain

$$\int_{-1}^0 w_n^{(s+1)}(x; \alpha, 1, r) dx = \frac{\beta_{n+1}(\alpha, r) - \beta_{n+1}(\alpha, r-s)}{s(n+1)}.$$

Finally, using (11) yields the desired equation. ■

We note that for $s = 1$, (20) becomes

$$\beta_{n+1}(\alpha, r) - \beta_{n+1}(\alpha, r-1) = (n+1)(r-1 | \alpha)_n,$$

which is extension of well-known identity for Bernoulli polynomials

$$B_{n+1}(r) - B_{n+1}(r-1) = (n+1)(r-1)^n.$$

Taking $s = r$ in (20), we have

$$\beta_{n+1}(\alpha, r) = \beta_{n+1}(\alpha) + (n+1) \sum_{k=0}^n S_2(n, k, r | \alpha) \frac{(-1)^k \langle r \rangle_{k+1}}{k+1}.$$

On the other hand, taking $s = r$ in (20) and using the relation [9, Eq. (5.4)]

$$\sum_{j=0}^{r-1} (j | \alpha)_m = \frac{1}{m+1} [\beta_{m+1}(\alpha, r) - \beta_{m+1}(\alpha)],$$

we derive the following corollary.

Corollary 5 *The sums of generalized falling factorials can be expressed as*

$$\sum_{j=0}^{r-1} (j | \alpha)_n = \sum_{k=0}^n S_2(n, k, r | \alpha) \frac{(-1)^k \langle r \rangle_{k+1}}{k+1},$$

where r is any integer > 0 .

Setting $s = \alpha$ in (20) and using the identity [9, Eq. (5.10)]

$$\beta_n(\alpha, z + \alpha) = \beta_n(\alpha, z) + \alpha n \beta_{n-1}(\alpha, z), \quad \alpha > 0,$$

gives the following corollary.

Corollary 6 *For every $\alpha \geq 0$,*

$$\beta_n(\alpha, r - \alpha) = \sum_{k=0}^n S_2(n, k, r | \alpha) \frac{(-1)^k \langle \alpha + 1 \rangle_k}{k + 1}. \quad (22)$$

The identity given in [33] can also be derived from (22) for $\alpha \rightarrow 0$. Moreover, when $r = 0$, (22) becomes

$$\beta_n(\alpha, -\alpha) = \sum_{k=0}^n S_2(n, k | \alpha) \frac{(-1)^k \langle \alpha + 1 \rangle_k}{k + 1}. \quad (23)$$

Taking $r = \alpha$ in (22) we have

$$\beta_n(\alpha) = \sum_{k=0}^n S_2(n, k, \alpha | \alpha) \frac{(-1)^k \langle \alpha + 1 \rangle_k}{k + 1}. \quad (24)$$

Let us return to (21) again. For $\alpha \rightarrow 0$, (21) can be written as

$$\begin{aligned} \lim_{\alpha \rightarrow 0} \sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-1}^0 w_n^{(s+1)}(x; \alpha, \beta, r) dx &= \frac{1}{st} \left[\frac{te^{rt}}{e^{\beta t} - 1} - \frac{te^{(r-\beta s)t}}{e^{\beta t} - 1} \right] \\ &= \frac{1}{s} \left[\sum_{n=0}^{\infty} \frac{B_{n+1}\left(\frac{r}{\beta}\right) - B_{n+1}\left(\frac{r}{\beta} - s\right)}{n + 1} \beta^{n+1} \right] \frac{t^n}{n!}. \end{aligned}$$

Since $S_2(n, k; 0, \beta, r) = W_{\beta, r}(n, k)$, we derive the values of Bernoulli polynomials at rational numbers in the following theorem.

Theorem 7 *For every $s \geq 0$ and $\beta \neq 0$, we have*

$$B_{n+1}\left(\frac{r}{\beta}\right) - B_{n+1}\left(\frac{r}{\beta} - s\right) = (n + 1) \sum_{k=0}^n W_{\beta, r}(n, k) \frac{(-1)^k \langle s \rangle_{k+1}}{\beta^{n+1-k} (k + 1)}. \quad (25)$$

We note that for $\beta = 1$ and $r = 0$ in (25) we have

$$B_{n+1}(-s) = B_{n+1} + (n + 1) \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \frac{(-1)^{k+1} \langle s \rangle_{k+1}}{k + 1}.$$

Replacing $-s$ with x in the above equation and using (13) gives the well-known relation between Bernoulli polynomials and falling factorial as [35, Eq. (15.39)]

$$B_{n+1}(x) = B_{n+1} + \sum_{k=0}^n \frac{(n + 1)}{k + 1} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_{k+1}.$$

For $\beta = 1$ in (25) we have

$$B_{n+1}(r) - B_{n+1}(r-s) = \sum_{k=0}^n \frac{(-1)^k (n+1)}{k+1} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r \langle s \rangle_{k+1}.$$

Thus, taking $s = r$ in the above equation gives a new relation between Bernoulli polynomials and rising factorial in the following corollary.

Corollary 8 *For every $r \geq 0$, we have*

$$B_{n+1}(r) = B_{n+1} + \sum_{k=0}^n \frac{(-1)^k (n+1)}{k+1} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r \langle r \rangle_{k+1}.$$

For the last consequences of Theorem 7, we deal with the Howard's identity [23, Eq. (4.3)]

$$\sum_{j=0}^{m-1} (r + \beta j)^n = \frac{\beta^n}{n+1} \left[B_{n+1} \left(m + \frac{r}{\beta} \right) - B_{n+1} \left(\frac{r}{\beta} \right) \right],$$

Replacing $-s$ with m in (25) and using (13) gives that the sums of powers of integers can be evaluated as in the following corollary.

Corollary 9 *Let n and m be non-negative integers with $m > 0$ and $\beta \neq 0$. Then, we have*

$$\sum_{j=0}^{m-1} (r + \beta j)^n = \sum_{k=0}^n \frac{\beta^{k-1}}{k+1} W_{\beta,r}(n, k) (m)_{k+1}.$$

4 Some examples of series and integrals evaluation

In this section, several examples for the evaluation of some series and integrals are given.

For the first example, let us take

$$f(x) = \cosh(x/\beta) = \sum_{k=0}^{\infty} \frac{x^{2k}}{\beta^{2k} (2k)!},$$

in (1). Then, we have

$$\begin{aligned} \left(\beta x^{1-\alpha/\beta} D \right)^n \left[x^{r/\beta} \left(e^{x/\beta} + e^{-x/\beta} \right) \right] &= 2 \sum_{k=0}^{\infty} \frac{1}{\beta^{2k} (2k)!} \left(\beta x^{1-\alpha/\beta} D \right)^n \left[x^{(2k\beta+r)/\beta} \right] \\ &= 2 x^{(r-n\alpha)/\beta} \sum_{k=0}^{\infty} \frac{(2k\beta+r \mid \alpha)_n}{\beta^{2k} (2k)!} x^{2k}. \end{aligned}$$

From (4), the left hand side can be written as

$$\begin{aligned}
\left(\beta x^{1-\alpha/\beta} D\right)^n \left[x^{r/\beta} \left(e^{x/\beta} + e^{-x/\beta}\right)\right] \\
= \left(\beta x^{1-\alpha/\beta} D\right)^n \left[x^{r/\beta} e^{x/\beta}\right] + \left(\beta x^{1-\alpha/\beta} D\right)^n \left[x^{r/\beta} e^{-x/\beta}\right] \\
= x^{(r-n\alpha)/\beta} e^{x/\beta} S_n(x) + x^{(r-n\alpha)/\beta} e^{-x/\beta} S_n(-x).
\end{aligned}$$

Thus, we have

$$2 \sum_{k=0}^{\infty} \frac{(2k\beta + r | \alpha)_n}{\beta^{2k} (2k)!} x^{2k} = e^{x/\beta} S_n(x) + e^{-x/\beta} S_n(-x).$$

Moreover, setting $x = 2\pi i\beta$ in the above equation gives

$$\sum_{k=0}^{\infty} (2k\beta + r | \alpha)_n \frac{(-1)^k (2\pi)^{2k}}{(2k)!} = \sum_{j=1}^{\lfloor n/2 \rfloor} S(n, 2j; \alpha, \beta, r) (-1)^j (2\pi\beta)^{2j}. \quad (26)$$

Similarly, taking $f(x) = \sinh x$ in (1) gives

$$2 \sum_{k=0}^{\infty} \frac{((2k+1)\beta + r | \alpha)_n}{\beta^{2k+1} (2k+1)!} x^{2k+1} = e^{x/\beta} S_n(x) - e^{-x/\beta} S_n(-x)$$

and

$$\sum_{k=0}^{\infty} ((2k+1)\beta + r | \alpha)_n \frac{(-1)^k (2\pi)^{2k}}{(2k+1)!} = \beta \sum_{j=1}^{\lfloor n/2 \rfloor} S(n, 2j+1; \alpha, \beta, r) (-1)^j (2\pi\beta)^{2j}. \quad (27)$$

Note that (26) and (27) are the generalization of the identities given in [32, Page 403].

Now, if we apply (1) to the both sides of the function

$$\frac{1}{(1-x)^{s+1}} = \sum_{k=0}^{\infty} \binom{s+k}{k} x^k$$

and use (12), we obtain (for $n, s = 0, 1, 2, \dots$)

$$\sum_{k=0}^{\infty} \binom{s+k}{k} (r+k\beta | \alpha)_n x^k = \frac{1}{(1-x)^{s+1}} w_n^{(s+1)} \left(\frac{x}{1-x}; \alpha, \beta, r \right), \quad (28)$$

which is the generalization of [25]

$$\sum_{k=0}^{\infty} (r+k\beta | \alpha)_n x^k = \frac{1}{(1-x)} w_n \left(\frac{x}{1-x}; \alpha, \beta, r \right). \quad (29)$$

For the next example, for every $s > 0$, if we take

$$f(x) = (1+x)^s = \sum_{k=0}^{\infty} \binom{s}{k} x^k$$

in (1) and use (14), we derive

$$\sum_{k=0}^{\infty} \binom{s}{k} (r+k\beta \mid \alpha)_n x^k = (1+x)^s w_n^{(-s)} \left(\frac{-x}{1+x}; \alpha, \beta, r \right),$$

which is the generalization of the identity in [7, Eq. 9].

For the last example of the evaluation of the series in closed forms, we deal with the digamma function $\psi(x)$. The digamma function $\psi(x)$ can be given by the Taylor series at $x = 1$, [2, Eq. 6.3.14]

$$\psi(x+1) = -\gamma + \sum_{k=1}^{\infty} \zeta(k+1) (-1)^{k+1} x^k, \quad |x| < 1, \quad (30)$$

where $\zeta(s)$ is the Riemann zeta function and γ is the Euler's constant. Taking (30) in (1) gives

$$\left(\beta x^{1-\alpha/\beta} D \right)^n \left[x^{r/\beta} \psi(x+1) \right] = x^{(r-n\alpha)/\beta} \sum_{k=0}^n S(n, k; \alpha, \beta, r) \beta^k x^k \psi^{(k)}(x+1), \quad (31)$$

where $\psi^{(m)}(x)$ is the polygamma function which can be written more compactly in terms of the Hurwitz zeta function $\zeta(s, x)$ as [2, Eq. 6.4.10]

$$\psi^{(m)}(x) = (-1)^{m+1} m! \zeta(m+1, x). \quad (32)$$

Here $m > 0$, and x is any complex number not equal to a negative integer. Using (30) in the left hand side and (32) in the right hand side of (31), we obtain the following theorem.

Theorem 10 For $|x| < 1$,

$$\begin{aligned} & \sum_{k=1}^{\infty} \zeta(k+1) (r+k\beta \mid \alpha)_n x^k \\ &= -(r \mid \alpha)_n (\psi(1-x) + \gamma) + \sum_{k=1}^n S(n, k; \alpha, \beta, r) k! \zeta(k+1, 1-x) (\beta x)^k. \end{aligned} \quad (33)$$

As a consequences of Theorem 10, the following sums are obtained:

For $(\alpha, \beta, r) = (0, \beta, r)$,

$$\sum_{k=1}^{\infty} \zeta(k+1) (r+k\beta)^n x^k = -r^n (\psi(1-x) + \gamma) + \sum_{k=1}^n W_{\beta, r}(n, k) k! \zeta(k+1, 1-x) (\beta x)^k.$$

For $(\alpha, \beta, r) = (0, 1, r)$,

$$\sum_{k=1}^{\infty} \zeta(k+1) (r+k)^n x^k = -r^n (\psi(1-x) + \gamma) + \sum_{k=1}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r k! \zeta(k+1, 1-x) x^k.$$

For $(\alpha, \beta, r) = (0, 1, 1)$, using the relation $\{n\}_k = \{n\}_{k-1} = \{n\}$,

$$\sum_{k=1}^{\infty} \zeta(k+1) (k+1)^n x^k = -(\psi(1-x) + \gamma) + \sum_{k=1}^n \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} k! \zeta(k+1, 1-x) x^k. \quad (34)$$

Finally, setting $x = 1/2$ in the above equation, using $\psi(1/2) = -\gamma - 2 \log 2$ and $\zeta(s, 1/2) = (2^s - 1) \zeta(s)$ yields

$$\sum_{k=2}^{\infty} \frac{\zeta(k) k^n}{2^k} = \log 2 + \sum_{k=1}^n \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} k! (1 - 2^{-k-1}) \zeta(k+1).$$

Therefore, some special cases of the above equation involving infinite sums of Riemann zeta function can be listed as

$$\begin{aligned} \sum_{k=2}^{\infty} \frac{\zeta(k)}{2^k} &= \log 2, \\ \sum_{k=2}^{\infty} \frac{\zeta(k) k}{2^k} &= \log 2 + \frac{3}{4} \zeta(2), \\ \sum_{k=2}^{\infty} \frac{\zeta(k) k^2}{2^k} &= \log 2 + \frac{9}{4} \zeta(2) + \frac{14}{8} \zeta(3). \end{aligned}$$

Note that similar result of (34) can be found in [7, Proposition 20].

Now, we want to prove the equation in (15) by using (1). Applying (1) to the both sides of the integral

$$\frac{1}{(1-x)^s} = \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-1} e^{-(1-x)t} dt$$

and using (12) we obtain

$$\frac{x^{(r-n\alpha)/\beta}}{(1-x)^{s+1}} w_n^{(s+1)} \left(\frac{x}{1-x}; \alpha, \beta, r \right) = \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-1} e^{-t} \left(\beta x^{1-\alpha/\beta} D \right)^n \left[x^{r/\beta} e^{xt} \right] dt.$$

At the same time, we have

$$\begin{aligned}
\left(\beta x^{1-\alpha/\beta} D\right)^n \left[x^{r/\beta} e^{xt}\right] &= \left(\beta x^{1-\alpha/\beta} D\right)^n \sum_{k=0}^{\infty} \frac{t^k}{k!} x^{k+r/\beta} \\
&= \sum_{k=0}^{\infty} \frac{t^k}{k!} \left(\beta x^{1-\alpha/\beta} D\right)^n x^{k+r/\beta} \\
&= x^{(r-n\alpha)/\beta} \sum_{k=0}^{\infty} \frac{t^k}{k!} (k\beta + r \mid \alpha)_n x^k \\
&= x^{(r-n\alpha)/\beta} e^{xt} S_n(x\beta t).
\end{aligned}$$

Thus, we have a different representation of (15) as

$$\frac{1}{(1-x)^{s+1}} w_n^{(s+1)} \left(\frac{x}{1-x}; \alpha, \beta, r \right) = \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-1} S_n(x\beta t) e^{-(1-x)t} dt.$$

Now, we want to add some examples for the evaluation of integrals. Before giving the examples we need to mention that in the rest of this section we use the well-known estimate for the gamma function:

$$|\Gamma(x + iy)| \sim \sqrt{2\pi} |y|^{x-\frac{1}{2}} e^{-x-\frac{\pi}{2}|y|},$$

($|y| \rightarrow \infty$) for any fixed real x . This explains the behavior of the gamma function on vertical lines $\{t = a + iz : -\infty < z < \infty, 0 < a < 1\}$.

For the first example, let us start from the Mellin integral representation [34, Formula 5.37]

$$\frac{1}{(1+x)^{s+1}} = \frac{1}{2\pi i \Gamma(s+1)} \int_{a-i\infty}^{a+i\infty} x^{-t} \Gamma(t) \Gamma(s+1-t) dt,$$

where $s \geq 0$, $0 < x < 1$. Apply (1) to the both sides of the above integral to obtain

$$\left(\beta x^{1-\alpha/\beta} D\right)^n \left[\frac{x^{r/\beta}}{(1+x)^{s+1}} \right] = \frac{x^{(r-n\alpha)/\beta}}{2\pi i \Gamma(s+1)} \int_{a-i\infty}^{a+i\infty} (r - \beta t \mid \alpha)_n x^{-t} \Gamma(t) \Gamma(s+1-t) dt.$$

From the left hand side of the above equation, we derive

$$\begin{aligned}
\left(\beta x^{1-\alpha/\beta} D\right)^n \left[\frac{x^{r/\beta}}{(1+x)^{s+1}} \right] &= \sum_{k=0}^{\infty} \binom{s+k}{k} (-1)^k \left(\beta x^{1-\alpha/\beta} D\right)^n x^{(k\beta+r)/\beta} \\
&= x^{(r-n\alpha)/\beta} \sum_{k=0}^{\infty} \binom{s+k}{k} (r + k\beta \mid \alpha)_n (-x)^k \\
&= \frac{x^{(r-n\alpha)/\beta}}{(1+x)^{s+1}} w_n^{(s+1)} \left(\frac{-x}{1+x}; \alpha, \beta, r \right).
\end{aligned}$$

Therefore, we have

$$\frac{1}{(1+x)^{s+1}} w_n^{(s+1)} \left(\frac{-x}{1+x}; \alpha, \beta, r \right) = \frac{1}{2\pi i \Gamma(s+1)} \int_{a-i\infty}^{a+i\infty} (r - \beta t | \alpha)_n x^{-t} \Gamma(t) \Gamma(s+1-t) dt, \quad (35)$$

Secondly, replace x by $(r + k\beta)x$, multiply both sides by $(r + k\beta | \alpha)_n$ and sum for $k = 0, 1, \dots$, in the integral:

$$e^{-x} = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} x^{-t} \Gamma(t) dt.$$

Then, we have

$$e^{-rx} \sum_{k=0}^{\infty} (r + k\beta | \alpha)_n (e^{-\beta x})^k = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} x^{-t} \sum_{k=0}^{\infty} \frac{(k\beta + r | \alpha)_n}{(k\beta + r)^t} \Gamma(t) dt.$$

From (29), the above integral becomes

$$\frac{e^{-rx}}{1 - e^{-\beta x}} w_n \left(\frac{e^{-\beta x}}{1 - e^{-\beta x}}; \alpha, \beta, r \right) = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} x^{-t} \sum_{k=0}^{\infty} \frac{(k\beta + r | \alpha)_n}{(k\beta + r)^t} \Gamma(t) dt.$$

The interesting part of the above integral, given in the following proposition, is appeared when $\alpha \rightarrow 0$.

Proposition 11 *For all $x > 0$, $\operatorname{Re}(\beta) > 0$, $\operatorname{Re}(r) > 0$, $n = 0, 1, 2, \dots$ and $a > n + 1$*

$$\frac{e^{-rx}}{1 - e^{-\beta x}} w_n \left(\frac{e^{-\beta x}}{1 - e^{-\beta x}}; 0, \beta, r \right) = \frac{\beta^n}{2\pi i} \int_{a-i\infty}^{a+i\infty} (\beta x)^{-t} \zeta \left(t - n, \frac{r}{\beta} \right) \Gamma(t) dt. \quad (36)$$

We note that for $\alpha \rightarrow 0$, generalized geometric polynomials become

$$w_n(x; 0, \beta, r) = \sum_{k=0}^n W_{\beta, r}(n, k) \beta^k k! x^k.$$

Moreover, for $\beta = 1$ and $n = 0$ in (36), we have the well-known inverse Mellin transformation of Hurwitz zeta function [3, Theorem 12.2]. Moreover, (35) and (36) are the generalization of identities in [7, Proposition 16].

References

- [1] C. Ahlbach, J. Usatine, N. Pippenger, Barred preferential arrangement, Electron. J. Combin. 20 (2) (2013) #P55.

- [2] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, New York Dover 1972.
- [3] T. M. Apostol, Introduction to analytic number theory, Undergraduate Texts in Mathematics, New York-Heidelberg Springer-Verlag (1976).
- [4] M. Benoumhani, On Whitney numbers of Dowling lattices, Discrete Math. 159 (1996) 13–33.
- [5] K. N. Boyadzhiev, A series transformation formula and related polynomials, Int. J. Math. Math. Sci. 23 (2005) 3849–3866.
- [6] K. N. Boyadzhiev, Exponential polynomials, Stirling numbers and evaluation of some gamma integrals, Abstr. Appl. Anal. 18 (2009) 1–18.
- [7] K. N. Boyadzhiev, A. Dil, Geometric polynomials: properties and applications to series with zeta values, Analysis Math. 42 (3) (2016) 203–224.
- [8] A. Z. Broder, The r -Stirling numbers, Discrete Math. 49 (1984) 241–259.
- [9] L. Carlitz, Degenerate Stirling, Bernoulli and Eulerian numbers, Util. Math. 15 (1979) 51–88.
- [10] M. Cenkci, F. T. Howard, Notes on degenerate numbers, Discrete Math 307 (2007) 2359–2375.
- [11] M. Cenkci, An explicit formula for generalized potential polynomials and its applications, Discrete Math, 309 (2009) 1498–1510.
- [12] L. Comtet, Advanced Combinatorics, D. Reidel Dordrecht 1974.
- [13] R. B. Corcino, Some theorems on generalized Stirling numbers, Ars Combin. 60 (2001) 273–286.
- [14] R. B. Corcino, C. B. Corcino, On generalized Bell polynomials, Discret Dyn Nat Soc. 2011 2011 21p.
- [15] R. B. Corcino, M. B. Montero, C. B. Corcino, On generalized Bell numbers for complex argument, Util. Math. 88 (2012) 267–279.
- [16] M.C. Dağlı, M. Can, On reciprocity formula of character Dedekind sums and the integral of products of Bernoulli polynomials, *J. Number Theory* **156** (2015) 105–124.
- [17] A. Dil, V. Kurt, Polynomials related to harmonic numbers and evaluation of harmonic number series I, Integers 12 (2012), A38.
- [18] A. Dil, V. Kurt, Polynomials related to harmonic numbers and evaluation of harmonic number series II, Appl. Anal. Discrete Math. 5 (2011), 212–229.
- [19] L. Euler, Institutiones Calculi Differentialis, Vol. II, Academie Imperialis Scientiarum Petropolitanae, 1755.

- [20] R. L. Graham, D. E. Knuth, O. Patashnik, Concrete Mathematics, Addison-Wesley Publ. Co. New York 1994.
- [21] F.T.Howard, Degenerate weighted Stirling numbers, Discrete Math. 57 (1985) 45–58.
- [22] F.T. Howard, Explicit formulas for degenerate Bernoulli numbers, Discrete Math. 162 (1996) 175–185.
- [23] F. T. Howard, Sums of powers of integers via generating functions, Fibonacci Quart. 34 (3) (1996) 244–256.
- [24] L. C. Hsu, P. J. S. Shiue, A unified approach to generalized Stirling numbers, Adv. in Appl. Math., 20 3 (1998) 366–384.
- [25] L. Kargin, R. B. Corcino, Generalization of Mellin derivative and its applications, Integral Transforms Spec. Funct. 27 (8) (2016) 620–631.
- [26] T. Kim, Jong Jin Seo, Degenerate Bernoulli Numbers and Polynomials of the Second Kind, International Journal of Mathematical Analysis 9 (26) 2015 1269-1278.
- [27] P. M. Knopf, The operator $(x\frac{d}{dx})^n$ and its application to series, Math. Mag. 76 (5) (2003) 364–371.
- [28] M. Merca, A new connection between r -Whitney numbers and Bernoulli polynomials. Integral Transforms Spec. Funct. 25 (12) (2014) 937–942.
- [29] M. Merca, A connection between Jacobi–Stirling numbers and Bernoulli polynomials, *J. Number Theory* 151 (2015) 223–229.
- [30] M. Merca, A note on the r -Whitney numbers of Dowling lattices, C. R. Acad. Sci. Paris, Ser. I 351 (2013) 649–655.
- [31] I. Mező, A new formula for the Bernoulli polynomials, Results in Mathematics, 58 (2010) 329–335.
- [32] I. Mező, J. L. Ramirez, Some identities for the r -Whitney numbers, Aequationes Mathematicae, 90 (2016) 393-406.
- [33] M. Mihoubi, M. Tiachachat, Some applications of the r -Whitney numbers, C.R. Acad. Sci. Paris Ser.I 352 (2014) 965–969.
- [34] F. Oberhettinger, Tables of Mellin Transforms, Springer Berlin. 1974.
- [35] J. Quaintance, H. W. Gould, Combinatorial Identities for Stirling Numbers: The unpublished notes of H. W. Gould, World Scientific Publishing Company, 2015.
- [36] A. Xu, Extensions of Spivey’s Bell number formula, Electron. J. Combin. 19 (2) (2012) p.8.

- [37] P. T. Young, Degenerate Bernoulli polynomials, generalized factorial sums, and their applications, *J. Number Theory*, 128 (4) (2008) 738-758.
- [38] P. T. Young, Congruences for degenerate number sequences, *Discrete Math*, 270, (1-3) (2003) 279-289.